



Three-dimensional Orographic Precipitation Enhancement Observed in OLYMPEX

R. A. Houze, Jr.,
Angela Rowe,
Lynn McMurdie,
Stacy Brodzik,
Megan Chaplin
Joe Zagrodnik,
Thomas Schuldt

See poster 244

Identification of Mesoscale Convective Precipitation in GPM Data

Robert Houze and Stacy Brodzik
PMM Science Team Meeting
San Diego, CA
October 16-20, 2017



Reviews of Geophysics

REVIEW ARTICLE

**The variable nature of convection in the tropics and subtropics:
A legacy of 16 years of the Tropical Rainfall Measuring
Mission satellite**

Robert A. Houze Jr.¹, Kristen L. Rasmussen², Manuel D. Zuluaga³, and Stella R. Brodzik¹


August 2015

The TRMM study identified
categories of convective and
stratiform echoes based on their
3D structure

Object	Echo Characteristic	Height	Area
Isolated Shallow Echoes	> min detectable	< 4 km	2 pixels
Deep Convective Cores	> 40 dBZ (<u>strong</u>)	> 10 km	N/A
	> 30 dBZ (<u>moderate</u>)	> 8 km	N/A
Wide Convective Cores	> 40 dBZ (<u>strong</u>)		> 1000 km ² at some altitude
	> 30 dBZ (<u>moderate</u>)		> 800 km ² at some altitude
Broad Stratiform Regions	Contiguous stratiform		> 50,000 km ² (<u>strong</u>)
			> 30,000 km ² (<u>moderate</u>)
			30-50,000 km ² (<u>filtered</u>)

An Example of the TRMM Results


Deep Convective Cores
(strong)




Frequencies of occurrence

See Houze et al. 2015 for
full low-latitude patterns of
these features

Wide Convective Cores
(strong)



Broad Stratiform
(strong)






GPM Goal:

Extend this work to higher latitudes

Problem:

- V4 classified a lot of Ku stratiform echo as convective
- V5 corrected much of this problem

Example of successful correction




October
Missouri

(20141009/1637
)


HUGE
improvement

But problems remain

Problem case: East Coast Front




October
NE U.S.




(20141022/1132)

Problem case: West Coast Front




December
NW coast of U.S.




(20141224/1202)

Problem case: North Pacific



February
North Pacific



(20150202/0839)

Solving this remaining problem is
critical to all applications that depend on
convective/stratiform classification


(latent heating, PSD, ...)

Complicating factors in identifying stratiform echo in GPM data:


- Melting level height varies
- Bright band is often >40 dBZ
- Melting level is not always a sharp peak in dBZ
- Key indicator of SF is a strong drop off in reflectivity above melting level

UW add-on to V5 C/S algorithm


Test Area




Convective Reflectivity Profile - 20140630/2147




Stratiform Reflectivity Profile - 20170801/2018




Reclassified Stratiform Reflectivity Profile - 20140630/2147




Reclassified Stratiform Reflectivity Profile - 20170801/2018






Problem case: East Coast Front



Problem case: West Coast Front



Problem case: North Pacific







February
North Pacific

(20150202/0839)





Object	Echo Characteristic	Height	Area
Isolated Shallow Echoes	> min detectable	< 4 km	2 pixels
Deep Convective Cores	> 40 dBZ (<u>strong</u>)	> 10 km	N/A
	> 30 dBZ (<u>moderate</u>)	> 8 km	N/A
Wide Convective Cores	> 40 dBZ (<u>strong</u>)		> 1000 km ² at some altitude
	> 30 dBZ (<u>moderate</u>)		> 800 km ² at some altitude
Broad Stratiform Regions	Contiguous stratiform		> 50,000 km ² (<u>strong</u>)
			> 30,000 km ² (<u>moderate</u>)
			30-50,000 km ² (<u>filtered</u>)

Comparison of v05 Counts NASA/UW (strong thresholds)







Comparison of v05 Counts

NASA/UW (moderate thresholds)




Percentage of WCCs reclassified by UW add-on (strong threshold cases)




PROBABILITY MAPS (BASED ON GPM-Ku v05, 2014-2017)


(a) DJF broad stratiform regions strong thresholds




(b) DJF deep convective cores strong thresholds



(c) DJF deep/wide convective cores strong thresholds




(d) DJF wide convective cores strong thresholds




PROBABILITY MAPS (BASED ON GPM-Ku v05uw2, 2014-2017)


(a) DJF broad stratiform regions strong thresholds (uw2)




(b) DJF deep convective cores strong thresholds (uw2)



(c) DJF deep/wide convective cores strong thresholds (uw2)




(d) DJF wide convective cores strong thresholds (uw2)




PROBABILITY MAPS (BASED ON GPM-Ku v05, 2014-2017)


(a) MAM broad stratiform regions strong thresholds




(b) MAM deep convective cores strong thresholds



(c) MAM deep/wide convective cores strong thresholds




(d) MAM wide convective cores strong thresholds




PROBABILITY MAPS (BASED ON GPM-Ku v05uw2, 2014-2017)


(a) MAM broad stratiform regions strong thresholds (uw2)




(b) MAM deep convective cores strong thresholds (uw2)



(c) MAM deep/wide convective cores strong thresholds (uw2)



(d) MAM wide convective cores strong thresholds (uw2)




Conclusions

- V5 performs better than V4 but still indicates a lot of stratiform as convective → overestimation of convective latent heating
- An algorithm based on the dropoff of reflectivity above the maximum dBZ eliminates most of the problem
- Over the North America sector, the problem is greatest in winter and spring, at higher latitudes, & over the ocean


Extra Slides

PROBABILITY MAPS (BASED ON GPM-Ku v05, 2014-2017)


(a) JJA broad stratiform regions strong thresholds




(b) JJA deep convective cores strong thresholds



(c) JJA deep/wide convective cores strong thresholds




(d) JJA wide convective cores strong thresholds




PROBABILITY MAPS (BASED ON GPM-Ku v05uw2, 2014-2017)


(a) JJA broad stratiform regions strong thresholds (uw2)




(b) JJA deep convective cores strong thresholds (uw2)



(c) JJA deep/wide convective cores strong thresholds (uw2)




(d) JJA wide convective cores strong thresholds (uw2)




PROBABILITY MAPS (BASED ON GPM-Ku v05, 2014-2016)


(a) SON broad stratiform regions strong thresholds




(b) SON deep convective cores strong thresholds



(c) SON deep/wide convective cores strong thresholds




(d) SON wide convective cores strong thresholds




PROBABILITY MAPS (BASED ON GPM-Ku v05uw2, 2014-2016)


(a) SON broad stratiform regions strong thresholds (uw2)



(b) SON deep convective cores strong thresholds (uw2)



(c) SON deep/wide convective cores strong thresholds (uw2)



(d) SON wide convective cores strong thresholds (uw2)

